

Affiliated Physician

1585 BARRINGTON RD, SUITE 603, BUILDING 2. HOFFMAN ESTATES, IL 60169, 847-310-1600

TRAUMA TO THE FOOT AND ANKLE

BELOW THE KNEE AMPUTATION VS LIMB SALVAGE. MAKE THE CORRECT CHOICE FOR YOUR PATIENT!



PILON FRACTURES, CALCANEAL FRACTURES, ANKLE FRACTURES, METATARSAL FRACTURES, TOE FRACTURES, and
SPRAINS OF THE FOOT AND ANKLE DIABETIC CHARCOT SURGICAL SPECIALIST

ACCEPT MERIDIAN, BCBS COMMUNITY AND PPO

Dr. Sutpal Singh, DPM, FACFAS: COMPREHENSIVE FOOT AND ANKLE INSTITUTE

Dr. Singh is one of a very few Foot and Ankle Surgeons in the United States who is Double Board Certified in Foot Surgery and Reconstructive Rearfoot and Ankle Surgery, *and* who has had 5 mini- fellowships (2 in Athens, Greece and 3 in the world renowned Ilizarov center in Kurgan, Russia), in leg lengthening, deformity correction and traumatology. His specialized training allows him to perform minimal incision surgery to correct deformities in the foot and ankle including bunion surgery and trauma surgery.


He has also performed Microvascular Surgery: Columbia Presbyterian Hospital in NY, and Plastic surgery flaps from Duke University. He has also done Neurosurgery course at Dellon Institute Johns Hopkins University

Affiliated Physician

1585 BARRINGTON RD, SUITE 603, BUILDING 2. HOFFMAN ESTATES, IL 60169, 847-310-1600

Diabetic Trauma and Limb Salvage

Severe Diabetic Ankle Fracture with peripheral vascular disease treated with a no incision surgical technique using an external fixator to correct tibial varus ankle fracture



Affiliated Physician

1585 BARRINGTON RD, SUITE 603, BUILDING 2. HOFFMAN ESTATES, IL 60169, 847-310-1600

Open Grade 3B Diabetic Ankle Fracture. Stabilized with the Ilizarov Method. Note complete consolidation of the ankle fracture. Complete healing in 10 weeks.



Affiliated Physician

1585 BARRINGTON RD, SUITE 603, BUILDING 2. HOFFMAN ESTATES, IL 60169, 847-310-1600

Severe Comminuted Ankle Fracture in a Diabetic

Severe Comminuted Pilon Fracture stabilized with the ~~Ilizarov~~ Ilizarov Method. The patient was allowed to ambulate as tolerated after one week. Note complete consolidation of the fracture.



Trust your patients only with a highly trained foot and ankle surgeon



Affiliated Physician

1585 BARRINGTON RD, SUITE 603, BUILDING 2. HOFFMAN ESTATES, IL 60169, 847-310-1600



Dr. Sutpal Singh

Foot and Ankle Surgeon

Trauma and Reconstructive Surgery

Diabetic Limb Salvage

Adult & Children:

Foot Ankle/Deformity

Specializing in

Minimal Incision Bunion Surgery